

Contractive Markov systems II

IVAN WERNER

Email: ivan_werner@pochta.ru

February 1, 2008

Abstract

We continue development of the theory of contractive Markov systems (CMSs) initiated in [16]. In this paper, we extend some results from [16], [18], [19] and [20] to the case of contractive Markov systems which have probabilities with a square summable variation by using some ideas of A. Johansson and A. Öberg [10]. In particular, we show that an irreducible CMS has a unique invariant Borel probability measure if the vertex sets form an open partition of the state space and the restrictions of the probability functions on their vertex sets have a square summable variation and are bounded away from zero.¹

1 Introduction

In [16], we introduced a theory of *contractive Markov systems (CMS)* which provides a unifying framework in so-called 'fractal' geometry. It extends the known theory of *iterated function systems (IFS) with place dependent probabilities*, which are contractive on average, [1][6] in a way that it also covers *graph directed constructions* of 'fractal' sets [12]. In particular, Markov chains associated with such systems naturally extend finite Markov chains and inherit some of their properties.

By a *Markov system* we mean a structure on a metric space which generates a Markov process on it and is given by a family

$$(K_{i(e)}, w_e, p_e)_{e \in E}$$

(see Fig. 1) where E is the set of edges of a finite directed (multi)graph (V, E, i, t) ($V := \{1, \dots, N\}$ is the set of vertices of the directed (multi)graph (we do not exclude the case $N = 1$), $i : E \rightarrow V$ is a map indicating the initial vertex

¹MSC 2000: 60J05, 37D35, 37A50, 37H99, 28A80. *Keywords:* contractive Markov systems (CMS), iterated function systems (IFS) with place-dependent probabilities, random systems with complete connections, g -measures, Markov chains, equilibrium states, fractals.

of each edge and $t : E \longrightarrow V$ is a map indicating the terminal vertex of each edge), $K_1, K_2, \dots, K_N$ is a partition of a metric space (K, d) into non-empty Borel subsets, $(w_e)_{e \in E}$ is a family of Borel measurable self-maps on the metric space such that $w_e(K_{i(e)}) \subset K_{t(e)}$ for all $e \in E$ and $(p_e)_{e \in E}$ is a family of Borel measurable probability functions on K (i.e. $p_e(x) \geq 0$ for all $e \in E$ and $\sum_{e \in E} p_e(x) = 1$ for all $x \in K$) (associated with the maps) such that each p_e is zero on the complement of $K_{i(e)}$.




Figure 1. A Markov system.

A Markov system is called *irreducible* or *aperiodic* iff its directed graph is irreducible or aperiodic respectively.

We call a Markov system $(K_{i(e)}, w_e, p_e)_{e \in E}$ *contractive* with an *average contracting rate* $0 < a < 1$ iff it satisfies the following *condition of contractiveness on average*

$$\sum_{e \in E} p_e(x) d(w_e x, w_e y) \leq a d(x, y) \text{ for all } x, y \in K_i, i = 1, \dots, N. \quad (1)$$

This condition was discovered by Richard Isaac in 1961 for the case $N = 1$ [7].

A Markov system $(K_{i(e)}, w_e, p_e)_{e \in E}$ determines a Markov operator U on the set of all bounded Borel measurable functions $\mathcal{L}^0(K)$ by

$$Uf := \sum_{e \in E} p_e f \circ w_e \text{ for all } f \in \mathcal{L}^0(K)$$

and its adjoint operator U^* on the set of all Borel probability measures $P(K)$ by

$$U^* \nu(f) := \int U(f) d\nu \text{ for all } f \in \mathcal{L}^0(K) \text{ and } \nu \in P(K).$$

Remark 1 Note that each map w_e and each probability p_e need to be defined only on the corresponding vertex set $K_{i(e)}$. This is sufficient for the condition (1) and the definition of U^* . For the definition of U , we can consider each w_e to be extended on the whole space K arbitrarily and each p_e to be extended on K by zero.

Also, the situation applies where each vertex set K_i has its own metric d_i . In this case, one can set

$$d(x, y) = \begin{cases} d_i(x, y) & \text{if } x, y \in K_i \\ \infty & \text{otherwise} \end{cases}$$

and use the convention $0 \times \infty = 0$.

We say a probability measure μ is an *invariant probability measure* of the Markov system iff it is a stationary initial distribution of the associated Markov process, i.e.

$$U^* \mu = \mu.$$

A Borel probability measure μ is called *attractive* measure of the CMS if

$$U^{*n} \nu \xrightarrow{w^*} \mu \text{ for all } \nu \in P(K),$$

where w^* means weak* convergence. Note that an attractive probability measure is a unique invariant probability measure of the CMS if U maps continuous functions on continuous functions. We will denote the space of all bounded continuous functions by $C_B(K)$.

The main result in [16] concerning the uniqueness of the invariant measure is the following (see Lemma 1 and Theorem 2 in [16]) (see also [17] for the case of constant probabilities $p_e|_{K_{i(e)}}$ and compact state space).

Theorem 1 Suppose $(K_{i(e)}, w_e, p_e)_{e \in E}$ is an irreducible CMS such that $K_1, K_2, \dots, K_N$ partition K into non-empty open subsets, each $p_e|_{K_{i(e)}}$ is Dini-continuous and there exists $\delta > 0$ such that $p_e|_{K_{i(e)}} \geq \delta$ for all $e \in E$. Then:

(i) The CMS has a unique invariant Borel probability measure μ , and $\mu(K_i) > 0$ for all $i = 1, \dots, N$.

(ii) If in addition the CMS is aperiodic, then

$$U^n f(x) \rightarrow \mu(f) \text{ for all } x \in K \text{ and } f \in C_B(K)$$

and the convergence is uniform on bounded subsets, i.e. μ is an attractive probability measure of the CMS.

A function $h : (X, d) \rightarrow \mathbb{R}$ is called *Dini-continuous* iff for some $c > 0$

$$\int_0^c \frac{\phi(t)}{t} dt < \infty$$

where ϕ is the modulus of uniform continuity of f , i.e.

$$\phi(t) := \sup\{|h(x) - h(y)| : d(x, y) \leq t, x, y \in X\}.$$

It is easily seen that the Dini-continuity is weaker than the Hölder and stronger than the uniform continuity. There is a well known characterization of the Dini-continuity.

Lemma 1 *Let $0 < c < 1$ and $b > 0$. A function h is Dini-continuous iff*

$$\sum_{n=0}^{\infty} \phi(bc^n) < \infty$$

where ϕ is the modulus of uniform continuity of h .

The proof is simple (e.g. see [16]).

Furthermore, with a Markov system $(K_{i(e)}, w_e, p_e)_{e \in E}$, which has an invariant Borel probability measure μ , also is associated a measure preserving transformation $S : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow (\Sigma, \mathcal{B}(\Sigma), M)$, which we call a *generalized Markov shift*, where $\Sigma := \{(\dots, \sigma_{-1}, \sigma_0, \sigma_1, \dots) : \sigma_i \in E \forall i \in \mathbb{Z}\}$ is the *code space* provided with the product topology, $\mathcal{B}(\Sigma)$ denotes Borel σ -algebra on Σ and M is a *generalized Markov measure* on $\mathcal{B}(\Sigma)$ given by

$$M(m[e_1, \dots, e_k]) := \int p_{e_1}(x) p_{e_2}(w_{e_1}x) \dots p_{e_k}(w_{e_{k-1}} \circ \dots \circ w_{e_1}x) d\mu(x)$$

for every cylinder set $m[e_1, \dots, e_k] := \{\sigma \in \Sigma : \sigma_m = e_1, \dots, \sigma_{m+k-1} = e_k\}$, $m \in \mathbb{Z}$, and S is the usual left shift map on Σ . It is easy to verify that S preserves measure M , since $U^*\mu = \mu$ (see [18]).

For a CMS, this two pictures, Markovian and dynamical, are related by a *coding map* $F : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow K$ which was constructed in [18]. It is defined, if K is a complete metric space and each $p_e|_{K_i(e)}$ is Dini-continuous and bounded away from zero, by

$$F(\sigma) := \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ for } M\text{-a.e. } \sigma \in \Sigma$$

where $x_i \in K_i$ for each $i = 1, \dots, N$ (the coding map does not depend on the choice of x_i 's modulo an M -zero set). We show in this paper that F is also well defined under a weaker continuity condition on the probability functions $p_e|_{K_i(e)}$.

Let denote by $\mathcal{F}$ the sub- σ -algebra on Σ generated by cylinder sets of the form $m[e_m, \dots, e_0]$, $m \leq 0$. Note that F is $\mathcal{F}$ - $\mathcal{B}(K)$ -measurable.

Example 1 Let $G := (V, E, i, t)$ be a finite irreducible directed (multi)graph. Let $\Sigma_G^- := \{(\dots, \sigma_{-1}, \sigma_0) : \sigma_m \in E \text{ and } t(\sigma_m) = i(\sigma_{m-1}) \forall m \in \mathbb{Z} \setminus \mathbb{N}\}$ (be

one-sided subshift of finite type associated with G) endowed with the metric $d(\sigma, \sigma') := 2^k$ where k is the smallest integer with $\sigma_i = \sigma'_i$ for all $k < i \leq 0$. Let g be a positive, Dini-continuous function on Σ_G such that

$$\sum_{y \in T^{-1}(\{x\})} g(y) = 1 \text{ for all } x \in \Sigma_G$$

where T is the right shift map on Σ_G^- . Set $K_i := \{\sigma \in \Sigma_G^- : t(\sigma_0) = i\}$ for every $i \in V$ and, for every $e \in E$,

$$w_e(\sigma) := (\dots, \sigma_{-1}, \sigma_0, e), \quad p_e(\sigma) := g(\dots, \sigma_{-1}, \sigma_0, e) \text{ for all } \sigma \in K_{i(e)}.$$

Obviously, maps $(w_e)_{e \in E}$ are contractions. Therefore, $(K_{i(e)}, w_e, p_e)_{e \in E}$ defines a CMS. An invariant probability measure of this CMS is called a g -measure. This notion was introduced by Keane [8] and further developed by Ledrappier [11], Walters [14], Berbee [2] and others.

Example 2 Let $\mathbb{R}^2$ be normed by $\|\cdot\|_1$. Let $K_1 := \{(x, y) \in \mathbb{R}^2 : y \geq 1\}$ and $K_2 := \{(x, y) \in \mathbb{R}^2 : y \leq -1\}$. Consider the following maps on $\mathbb{R}^2$:

$$\begin{aligned} w_1 \begin{pmatrix} x \\ y \end{pmatrix} &:= \begin{pmatrix} -\frac{1}{2}x - 1 \\ -\frac{3}{2}y + \frac{1}{2} \end{pmatrix}, \quad w_2 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} -\frac{3}{2}x + 1 \\ \frac{1}{4}y + \frac{3}{4} \end{pmatrix}, \\ w_3 \begin{pmatrix} x \\ y \end{pmatrix} &:= \begin{pmatrix} -\frac{1}{2}|x| + 1 \\ -\frac{3}{2}y - \frac{1}{2} \end{pmatrix}, \quad \text{and } w_4 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} \frac{3}{2}|x| - 1 \\ -\frac{1}{4}y + \frac{3}{4} \end{pmatrix} \end{aligned}$$

with probability functions

$$\begin{aligned} p_1 \begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15} \sin^2 \|(x, y)\|_1 + \frac{53}{105} \right) 1_{K_1}(x, y), \\ p_2 \begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15} \cos^2 \|(x, y)\|_1 + \frac{3}{7} \right) 1_{K_1}(x, y), \\ p_3 \begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15} \sin^2 \|(x, y)\|_1 + \frac{53}{105} \right) 1_{K_2}(x, y) \text{ and} \\ p_4 \begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15} \cos^2 \|(x, y)\|_1 + \frac{3}{7} \right) 1_{K_2}(x, y). \end{aligned}$$

A simple calculation shows that $(K_{i(e)}, w_e, p_e)_{e \in \{1, 2, 3, 4\}}$, where $i(1) = i(2) = 1$ and $i(3) = i(4) = 2$, defines a CMS with an average contracting rate 209/210 on $K_1 \cup K_2$. Note that none of the maps are contractive (by Theorem 1, it has a unique (attractive) invariant Borel probability measure).

2 Main Part

Let $\mathcal{M} := (K_{i(e)}, w_e, p_e)_{e \in E}$ be a contractive Markov system with an average contracting rate $0 < a < 1$ and an invariant Borel probability measure μ .

We assume that: (K, d) is a metric space in which sets of finite diameter are relatively compact; the family $K_1, \dots, K_N$ partitions K into non-empty open subsets; each probability function $p_e|_{K_{i(e)}}$ is uniformly continuous and bounded away from zero by $\delta > 0$; the set of edges E is finite and the map $i : E \rightarrow V$ is surjective. Note that the assumption on the metric space implies that it is locally compact separable and complete.

In contrast to the papers [16], [18], [19] and [20], here we will prove some similar results under a weaker continuity condition on the probability functions $p_e|_{K_{i(e)}}$ than the Dini-continuity. In Subsection 2.1, we show that the coding map is well defined under this condition. In Subsection 2.2, using the coding map, we set up a thermodynamic formalism for CMSs, which is beyond the scope of the existing theory. Finally, using the developed thermodynamic formalism, we show in Subsection 2.3 that the irreducible CMS $\mathcal{M}$ with the probabilities $p_e|_{K_{i(e)}}$ satisfying the following continuity condition has a unique invariant Borel probability measure, which also can be obtained empirically.

Definition 1 We say that a function $h : K \rightarrow \mathbb{R}$ has a *square summable variation* iff for any $c > 0$

$$\int_0^c \frac{\phi^2(t)}{t} dt < \infty$$

where ϕ is the *modulus of uniform continuity* of h , i.e.

$$\phi(t) := \sup\{|h(x) - h(y)| : d(x, y) \leq t, x, y \in X\}$$

or equivalently, for any $b > 0$ and $0 < c < 1$,

$$\sum_{n=0}^{\infty} \phi^2(bc^n) < \infty,$$

which is obviously a weaker condition than the Dini-continuity and stronger than the uniform continuity.

This condition was introduced by A. Johansson and A. Öberg in [10], where they proved the uniqueness of the invariant probability measure for some iterated function systems with place-dependent probabilities satisfying this condition on a compact metric space. Shortly after that, it was announced by Berger, Hoffman and Sidoravicius [3] that the condition of the square summability of variation is tight, in the sense that for any $\epsilon > 0$ there exists a g -function with a summable variation to the power $2 + \epsilon$ such that the corresponding CMS as in Example 1 has several invariant Borel probability measures.

Example 3 Let $\alpha, \delta > 0$ such that $\alpha + \delta < 1$ and $0 < c < 1/e$. Set

$$p(x) = \begin{cases} \frac{\alpha}{\log \frac{1}{x}} + \delta & \text{if } 0 < x \leq \frac{1}{e} \\ \delta & \text{if } x = 0. \end{cases}$$

Let ϕ be the modulus of uniform continuity of p . Then

$$\phi(c^n) = \frac{\alpha}{n \log \frac{1}{c}} \text{ for all } n \in \mathbb{N}.$$

Hence p is not Dini-continuous, but obviously it has a square summable variation. See [10] for a discussion of the relation between the Johansson-Öberg condition and the Berbee condition [2].

Let $\Sigma^+ := E^{\mathbb{N}}$ endowed with the product topology of the discrete topologies. For $x \in K$, let P_x denote the Borel probability measure on Σ^+ given by

$$P_x([e_1, \dots, e_k]) := p_{e_1}(x) p_{e_2}(w_{e_1} x) \dots p_{e_k}(w_{e_{k-1}} \circ \dots \circ w_{e_1} x)$$

for every cylinder set $[e_1, \dots, e_k] \subset \Sigma^+$. Obviously, P_x represents the Markov process with the Dirac initial distribution concentrated at the point x .

All results for CMSs with probabilities with a square summable variation which we intend to prove in this paper follow from the next lemma. It generalizes Lemma 3 in [6] and Lemma 2.3 in [19]. For the proof of it we use the methodology of Johansson and Öberg [10].

Lemma 2 *Suppose $\mathcal{M}$ is a CMS such that each probability function $p_e|_{K_{i(e)}}$ has a square summable variation. Then P_y is absolutely continuous with respect to P_x for all $x, y \in K_i$ and $i = 1, \dots, N$.*

Proof. Let $\mathcal{A}_n$ be the σ -algebra on Σ^+ generated by the cylinders of the form $[e_1, \dots, e_n]$ for all $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, let X_n be a function of Σ^+ given by

$$X_n(\sigma) := \sum_{e_1, \dots, e_n} \frac{P_y([e_1, \dots, e_n])}{P_x([e_1, \dots, e_n])} 1_{[e_1, \dots, e_n]}(\sigma) \text{ for all } \sigma \in \Sigma^+$$

with the convention that $0/0 = 0$. Obviously, by the definition of P_x , we can restrict the summation in the definition of X_n on paths $(e_1, \dots, e_n)^*$ of the digraph.

Now, observe that, for all $m \leq n$ and $C_m \in \mathcal{A}_m$,

$$\int_{C_m} X_n dP_x = \sum_{C_n \subset C_m} P_y(C_n) = P_y(C_m) = \int_{C_m} X_m dP_x. \quad (2)$$

Hence, $(X_n, \mathcal{A}_n)_{n \in \mathbb{N}}$ is a P_x -martingale. If X_n is uniformly bounded, then there exists $X \in \mathcal{L}^1(P_x)$ such that $X_n \rightarrow X$ P_x -a.e. and in L^1 sense, and $E_{P_x}(X|\mathcal{A}_m) = X_m$ P_x -a.e. for all m (see [4]). Then, by (2),

$$\int_{C_m} X dP_x = \int_{C_m} X_m dP_x = P_y(C_m) \text{ for all } C_m \in \mathcal{A}_m.$$

It means that the Borel probability measures XP_x and P_y agree on all cylinder subsets of Σ^+ , and therefore, are equal. This implies the claim. So, it remains to show that the martingale $(X_n, \mathcal{A}_n)_{n \in \mathbb{N}}$ is uniformly integrable, i.e.

$$\sup_n \int_{X_n > K} X_n dP_x \rightarrow 0 \text{ as } K \rightarrow \infty.$$

By (2), it is equivalent to show that

$$\sup_n P_y(X_n > K) \rightarrow 0 \text{ as } K \rightarrow \infty.$$

The latter is equivalent to

$$\sup_n P_y(\log X_n > K) \rightarrow 0 \text{ as } K \rightarrow \infty.$$

Let us use the following abbreviation:

$$p_i^x(\sigma) := p_{\sigma_i}(w_{\sigma_{i-1}} \circ \dots \circ w_{\sigma_1} x)$$

for $i \geq 2$ and $p_1^x(\sigma) := p_{\sigma_1}(x)$ for all $\sigma \in \Sigma^+$. Then, since $\log x \leq x - 1$,

$$\begin{aligned} \log X_n &= \sum_{(e_1, \dots, e_n)^*} \log \frac{p_1^y \dots p_n^y}{p_1^x \dots p_n^x} 1_{[e_1, \dots, e_n]} \\ &= \sum_{(e_1, \dots, e_n)^*} \sum_{i=1}^n \log \frac{p_i^y}{p_i^x} 1_{[e_1, \dots, e_n]} \\ &\leq \sum_{(e_1, \dots, e_n)^*} \sum_{i=1}^n \frac{p_i^y - p_i^x}{p_i^x} 1_{[e_1, \dots, e_n]}. \end{aligned}$$

Now, observe that

$$\frac{p_i^y - p_i^x}{p_i^x} = \frac{p_i^y - p_i^x}{p_i^y} + \frac{(p_i^y - p_i^x)^2}{p_i^x p_i^y}.$$

Therefore,

$$\log X_n \leq Y_n + Z_n, \tag{3}$$

where

$$Y_n := \sum_{(e_1, \dots, e_n)^*} \sum_{i=1}^n \frac{p_i^y - p_i^x}{p_i^y} 1_{[e_1, \dots, e_n]}$$

and

$$Z_n := \frac{1}{\delta^2} \sum_{(e_1, \dots, e_n)^*} \sum_{i=1}^n (p_i^y - p_i^x)^2 1_{[e_1, \dots, e_n]}.$$

Furthermore, observe that

$$Y_{i+1} - Y_i = \sum_{(e_1, \dots, e_n)^*} \frac{p_{n+1}^y - p_{n+1}^x}{p_{n+1}^y} 1_{[e_1, \dots, e_n]} \text{ for all } i \geq 1.$$

and

$$\begin{aligned} & \int_{1[e_1, \dots, e_n]} (Y_{n+1} - Y_n) dP_y \\ &= \sum_{e_{n+1}} \frac{p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} y) - p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} x)}{p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} y)} \\ & \quad \times p_{e_1}(y) \dots p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_1} y) p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} y) \\ &= \sum_{e_{n+1}} (p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} y) - p_{e_{n+1}}(w_{e_n} \circ \dots \circ w_{e_1} x)) \\ & \quad \times p_{e_1}(y) \dots p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_1} y) \\ &= 0. \end{aligned}$$

Hence, $(Y_n, \mathcal{A}_n)_{n \in \mathbb{N}}$ is a P_y -martingale. Therefore, $Y_n - Y_{n-1}$, $Y_{n-1} - Y_{n-2}, \dots$, $Y_2 - Y_1$, Y_1 are orthogonal in $\mathcal{L}^2(P_y)$. By the Pythagoras equality, this implies that

$$\begin{aligned} \int Y_n^2 dP_y &= \int \left(\sum_{i=2}^n (Y_i - Y_{i-1}) + Y_1 \right)^2 dP_y \\ &= \sum_{i=2}^n \int (Y_i - Y_{i-1})^2 dP_y + \int Y_1^2 dP_y \\ &\leq \sum_{i=1}^n \int \frac{(p_i^y - p_i^x)^2}{\delta^2} dP_y \\ &\leq \frac{1}{\delta^2} \sum_{i=1}^n \int [p_{\sigma_i}(w_{\sigma_{i-1}} \circ \dots \circ w_{\sigma_1} y) - p_{\sigma_i}(w_{\sigma_{i-1}} \circ \dots \circ w_{\sigma_1} x)]^2 dP_y. \end{aligned}$$

Let

$$A_i := \left\{ \sigma \in \Sigma^+ : d(w_{\sigma_i} \circ \dots \circ w_{\sigma_1} y, w_{\sigma_i} \circ \dots \circ w_{\sigma_1} x) > a^{\frac{i}{2}} d(x, y) \right\}$$

for $i \in \mathbb{N}$. By the contractiveness on average condition,

$$\int d(w_{\sigma_i} \circ \dots \circ w_{\sigma_1} y, w_{\sigma_i} \circ \dots \circ w_{\sigma_1} x) dP_y \leq a^i d(x, y) \text{ for all } i.$$

Hence, by the Markov inequality,

$$P_y(A_i) \leq a^{\frac{i}{2}} \text{ for all } i.$$

Therefore,

$$\begin{aligned} \int Y_n^2 dP_y &\leq \frac{1}{\delta^2} \sum_{i=1}^n \left(P_y(A_i) + \phi^2 \left(a^{\frac{i}{2}} d(x, y) \right) \right) \\ &\leq \frac{1}{\delta^2} \left(\sum_{i=1}^{\infty} a^{\frac{i}{2}} + \sum_{i=1}^{\infty} \phi^2 \left(a^{\frac{i}{2}} d(x, y) \right) \right) \end{aligned}$$

for all $n \in \mathbb{N}$. Hence

$$\sup_n P_y(Y_n > K) \rightarrow 0 \text{ as } K \rightarrow \infty.$$

Analogously,

$$\sup_n P_y(Z_n > K) \rightarrow 0 \text{ as } K \rightarrow \infty.$$

Since, by (3),

$$\{\log X_n > K\} \subset \left\{ Y_n > \frac{K}{2} \right\} \cup \left\{ Z_n > \frac{K}{2} \right\},$$

we conclude that

$$\sup_n P_y(\log X_n > K) \rightarrow 0 \text{ as } K \rightarrow \infty,$$

as desired. $\square$

2.1 Coding map

We shall denote by d' the metric on Σ defined by $d'(\sigma, \sigma') := (1/2)^k$ where k is the largest integer with $\sigma_i = \sigma'_i$ for all $|i| < k$. Denote by $\mathcal{A}$ the finite σ -algebra generated by the zero time partition $\{0[e] : e \in E\}$ of Σ , and define, for each integer $m \leq 1$,

$$\mathcal{A}_m := \bigvee_{i=m}^{+\infty} S^{-i} \mathcal{A},$$

which is the smallest σ -algebra containing all finite σ -algebras $\bigvee_{i=m}^n S^{-i} \mathcal{A}$, $n \geq m$. Let $x \in K$. For each integer $m \leq 1$, let P_x^m be the probability measure on the σ -algebra $\mathcal{A}_m$ given by

$$P_x^m(m[e_m, \dots, e_n]) = p_{e_m}(x) p_{e_{m+1}}(w_{e_m}(x)) \dots p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_m}(x))$$

for all cylinder sets $m[e_m, \dots, e_n]$, $n \geq m$.

Lemma 3 *Let $m \leq 1$ and $A \in \mathcal{A}_m$. Then $x \mapsto P_x^m(A)$ is a Borel measurable function on K .*

The proof is simple (see e.g. [16]).

Definition 2 Let $\nu \in P(K)$. We call a probability measure $\Phi_m(\nu)$ on $(\Sigma, \mathcal{A}_m)$ given by

$$\Phi_m(\nu)(A) := \int P_x^m(A) d\nu(x), \quad A \in \mathcal{A}_m,$$

the m -th lift of ν .

Definition 3 Set

$$\mathcal{C}(B) := \left\{ (A_m)_{m=0}^{-\infty} : A_m \in \mathcal{A}_m \quad \forall m \text{ and } B \subset \bigcup_{m=0}^{-\infty} A_m \right\}$$

for $B \subset \Sigma$. Let $\nu \in P(K)$. We call a set function given by

$$\Phi(\nu)(B) := \inf \left\{ \sum_{m=0}^{-\infty} \Phi_m(\nu)(A_m) : (A_m)_{m \leq 0} \in \mathcal{C}(B) \right\}, \quad B \subset \Sigma,$$

the lift of ν .

Lemma 4 Let $\nu, \lambda \in P(K)$. Then

(i) $\Phi(\nu)$ is an outer measure on Σ .

(ii) If $\Phi_m(\nu) \ll \Phi_m(\lambda)$ for all $m \leq 0$, then for all $\epsilon > 0$ there exists $\alpha > 0$ such that

$$\Phi(\lambda)(B) < \alpha \Rightarrow \Phi(\nu)(B) < \epsilon \text{ for all } B \subset \Sigma.$$

For the proof see [18].

Note that $M = \Phi(\mu)$ (see [18]), where M is the generalized Markov measure associated with the CMS $\mathcal{M}$ and its invariant measure μ .

Fix $x_i \in K_i$ for each $i \in \{1, \dots, N\}$ and set

$$P_{x_1 \dots x_N}^m := \Phi_m \left(\frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right) \text{ and } P_{x_1 \dots x_N} := \Phi \left(\frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$

for every $m \in \mathbb{Z} \setminus \mathbb{N}$, where δ_x denotes the Dirac probability measure concentrated at x , i.e.

$$P_{x_1 \dots x_N}^m(m[e_m, \dots, e_n]) = \frac{1}{N} p_{e_m}(x_{i(e_m)}) p_{e_{m+1}}(w_{e_m} x_{i(e_m)}) \dots p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_m} x_{i(e_m)})$$

for every cylinder set $m[e_m, \dots, e_n]$.

Theorem 2 Let $x_i, y_i \in K_i$ for each $1 \leq i \leq N$. Then the following hold.

(i)

$$\lim_{m \rightarrow -\infty} d(w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(y_{i(\sigma_m)})) = 0$$

$P_{x_1 \dots x_N}$ -a.e..
(ii)

$$F_{x_1 \dots x_N} := \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(x_{i(\sigma_m)}) \text{ exists } P_{x_1 \dots x_N}\text{-a.e.},$$

and by (i) $F_{x_1 \dots x_N} = F_{y_1 \dots y_N}$ $P_{x_1 \dots x_N}$ -a.e..

(iii) There exists a sequence of closed subsets $Q_1 \subset Q_2 \subset \dots \subset \Sigma$ with $\sum_{k=1}^{\infty} P_{x_1 \dots x_N}(\Sigma \setminus Q_k) < \infty$ such that $F_{x_1 \dots x_N}|_{Q_k}$ is locally Hölder-continuous with the same Hölder-constants for all $k \in \mathbb{N}$, i.e. there exist $\alpha, C > 0$ such that for every k there exists $\delta_k > 0$ such that

$$\sigma, \sigma' \in Q_k \text{ with } d'(\sigma, \sigma') \leq \delta_k \Rightarrow d(F_{x_1 \dots x_N}(\sigma), F_{x_1 \dots x_N}(\sigma')) \leq C d'(\sigma, \sigma')^\alpha.$$

For the proof see [18].

Lemma 5 Suppose $(K_{i(e)}, w_e, p_e)_{e \in E}$ is a CMS such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Let $\nu_1 \in P(K)$ such that $\nu_1(K_i) > 0$ for all $i = 1, \dots, N$. Then $\Phi(\nu_2)$ is absolutely continuous with respect to $\Phi(\nu_1)$ for all $\nu_2 \in P(K)$.

Proof. By Lemma 4 (ii), it is sufficient to show that $\Phi_m(\nu_2)$ is absolutely continuous with respect to $\Phi_m(\nu_1)$ for all $m \leq 0$. Let $A \in \mathcal{A}_m$ such that $\Phi_m(\nu_1)(A) = 0$. Then for all $i = 1, \dots, N$ there exists $x_i \in K_i$ such that $P_{x_i}^m(A) = 0$. Hence, by the hypothesis, $P_x^m(A) = 0$ for all $x \in K$. Therefore,

$$\Phi_m(\nu_1)(A) = \int P_x^m(A) d\nu_1(x) = 0 \text{ for all } \nu_1 \in P(K),$$

as desired. $\square$

Corollary 1 Suppose $(K_{i(e)}, w_e, p_e)_{e \in E}$ is a CMS with an invariant Borel probability measure μ such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Let $x_i, y_i \in K_i$ for each $1 \leq i \leq N$. Then the following hold.

(i)

$$\lim_{m \rightarrow -\infty} d(w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(y_{i(\sigma_m)})) = 0 \text{ } M\text{-a.e.},$$

(ii)

$$F_{x_1 \dots x_N} := \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m}(x_{i(\sigma_m)}) \text{ exists } M\text{-a.e.},$$

and by (i) $F_{x_1 \dots x_N} = F_{y_1 \dots y_N}$ M -a.e..

(iii) There exists a sequence of closed subsets $Q_1 \subset Q_2 \subset \dots \subset \Sigma$ with $\lim_{k \rightarrow \infty} M(\Sigma \setminus Q_k) = 0$ such that $F_{x_1 \dots x_N}|_{Q_k}$ is locally Hölder-continuous with the

same Hölder-constants for all $k \in \mathbb{N}$, i.e. there exist $\alpha, C > 0$ such that for every k there exists $\delta_k > 0$ such that

$$\sigma, \sigma' \in Q_k \text{ with } d'(\sigma, \sigma') \leq \delta_k \Rightarrow d(F_{x_1 \dots x_N}(\sigma), F_{x_1 \dots x_N}(\sigma')) \leq C d'(\sigma, \sigma')^\alpha,$$

where M is the generalized Markov measure associated with the CMS and μ .

Proof. Set

$$\nu_1 := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}.$$

Since $M = \phi(\mu)$, the claim follows by Theorem 2 and Lemma 5. $\square$

2.2 The generalized Markov measure as an equilibrium state

Now, we are going to set up a thermodynamic formalism for the CMS $\mathcal{M}$. We will construct an energy function u with respect to which the generalized Markov measure M will turn out to be a unique equilibrium state if the probabilities $p_e|_{K_{i(e)}}$ of the CMS have a square summable variation. This extends the results from [20].

Definition 4 Let X be a metric space and T a continuous transformation on it. Denote by $P(X)$ the set of all Borel probability measures on X and by $P_T(X)$ the set of all T -invariant Borel probability measures on X . We call a Borel measurable function $h : X \rightarrow [-\infty, 0]$ an *energy function*. Suppose that T has a finite topological entropy, i.e. $\sup_{\Theta \in P_T(X)} h_\Theta(T) < \infty$, where $h_\Theta(T)$ is the Kolmogorov-Sinai entropy of T with respect to measure Θ . We call

$$P(h) = \sup_{\Theta \in P_T(X)} (h_\Theta(T) + \Theta(h))$$

the *pressure* of h . We call $\Lambda \in P_T(X)$ an *equilibrium state* for h iff

$$h_\Lambda(T) + \Lambda(h) = P(h).$$

Let's denote the set of all equilibrium states for h by $ES(h)$.

The construction of the function u goes through a definition of an appropriate shift invariant subset of Σ on which the energy function shall be finite. This subset is exactly that on which the coding map is defined. It means that the existence of an equilibrium state is closely related with the existence of the coding map.

Definition 5 Let

$$\Sigma_G := \{\sigma \in \Sigma : t(\sigma_j) = i(\sigma_{j+1}) \ \forall j \in \mathbb{Z}\},$$

$$D := \{\sigma \in \Sigma_G : \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ exists}\}$$

and

$$Y := \bigcap_{i=-\infty}^{\infty} S^i(D).$$

Now, set

$$u(\sigma) = \begin{cases} \log p_{\sigma_1} \circ F(\sigma) & \text{if } \sigma \in Y \\ -\infty & \text{if } \sigma \in \Sigma \setminus Y. \end{cases} \quad (4)$$

Lemma 6 (i) $F(\sigma)$ is defined for all $\sigma \in Y$ and Y is a shift invariant subset of Σ_G .

(ii) If P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$, then $M(Y) = 1$.

(iii) If P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$, and the CMS has an invariant probability measure μ such that $\mu(K_{i(e)}) > 0$ for all $e \in E$. Then Y is dense in Σ_G .

Proof. (i) is clear, by the definitions of F and Y . Note that the condition of the contractiveness on average and the boundedness away from zero of the functions $p_e|_{K_{i(e)}}$ imply that each map $w_e|_{K_{i(e)}}$ is continuous (Lipschitz). Therefore, $S(D) \subset D$. By Corollary 1, $M(D) = 1$ if P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. This implies (ii). If in addition $\mu(K_{i(e)}) > 0$ for all $e \in E$, then $M(O) > 0$ for every open $O \subset \Sigma_G$. This implies (iii). $\square$

Remark 2 (i) Note that Y and F depend on the choice of x_i 's. By Corollary 1 (ii), Y changes only modulo M -zero set by a different choice of x_i 's if P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$.

(ii) If all maps $w_e|_{K_{i(e)}}$ are contractive, then $Y = \Sigma_G$ and $F|_{\Sigma_G}$ is Hölder-continuous (easy to check).

Remark 3 As far as the author is aware, the rigorous mathematical theory of equilibrium states is developed only for upper semicontinuous energy functions h (see e.g. [9]). This condition insures that the convex space $ES(h)$ is non-empty and compact in the weak* topology. However, as the next example shows, u is not upper semicontinuous in general. Therefore, even the existence of an equilibrium state for u is not guaranteed by the existing theory.

Example 4 Let $(K, d) = (\mathbb{R}, |\cdot|)$. Consider two maps

$$w_0(x) := \frac{1}{2}x, \quad w_1(x) := 2x \text{ for all } x \in \mathbb{R}$$

with probability functions

$$p_0(x) := \frac{1}{6} \sin^2 x + \frac{17}{24}, \quad p_1(x) := \frac{1}{6} \cos^2 x + \frac{1}{8} \text{ for all } x \in \mathbb{R}.$$

Then a simple calculation shows that $(\mathbb{R}, w_e, p_e)_{e \in \{0,1\}}$ defines a CMS with an average contracting rate $45/48$. In this case, $\Sigma_G = \{0, 1\}^{\mathbb{Z}}$. If we take $x = 0$ for the definition of Y , then, obviously, $Y = \Sigma_G$. Now, let $x \neq 0$. Let $N_{0n}(\sigma)$ and $N_{1n}(\sigma)$ be the numbers of zeros and ones in $(\sigma_{-n}, \dots, \sigma_0)$ respectively for every $\sigma \in \Sigma_G$. Then, obviously, $\sigma \notin Y$ if $(N_{1n}(\sigma) - N_{0n}(\sigma)) \rightarrow \infty$. Hence, $Y \neq \Sigma_G$ and, by Lemma 6 (iii), Y is a dense shift invariant subset of Σ_G . Since Y is not closed, u is not upper semicontinuous.

Also, it is not difficult to see that in a general case there is no hope to find x_i such that u becomes upper semicontinuous, e.g. change w_1 to $w_1(x) = 2x + 1$, then, for any choice of x for the definition of Y , $Y \neq \Sigma_G$.

Proposition 1 *There exists an invariant Borel probability measure μ of the CMS $\mathcal{M}$ such that*

$$\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty \text{ for all } x_i \in K_i, i = 1, \dots, N.$$

Proof. Fix $x_i \in K_i$ for all $i = 1, \dots, N$ and set

$$f(x) := \sum_{i=1}^N 1_{K_i}(x) d(x, x_i) \text{ for all } x \in K.$$

Let $C := \max_{e \in E} d(w_e x_{i(e)}, x_{t(e)})$. Then the contractiveness on average condition (1) implies that

$$U^k f(x_i) \leq \frac{C}{1-a} \text{ for all } k \in \mathbb{N} \text{ and } i = 1, \dots, N$$

(see the proof of Theorem 1 in [16]).

Now, set

$$U_n := \frac{1}{n} \sum_{k=1}^n U^k \text{ for all } n \in \mathbb{N}.$$

By Theorem 1 (i) from [16], the sequence $(U^{*k} \delta_{x_i})_{k \in \mathbb{N}}$ is tight. Hence, the sequence $(U_n^* \delta_{x_i})_{n \in \mathbb{N}}$ is tight also. So, it has a subsequence, say $(U_{n_m}^* \delta_{x_i})_{m \in \mathbb{N}}$, which converges weakly* to a Borel probability measure, say μ . By the setup of $\mathcal{M}$, the operator U has the Feller property. This implies that $U^* \mu = \mu$.

Let $R > 0$ and $f_R := \min\{f, R\}$. Then

$$U_{n_m}^* \delta_{x_i}(f_R) \rightarrow \mu(f_R) \text{ as } m \rightarrow \infty.$$

On the other hand

$$U_{n_m}^* \delta_{x_i}(f_R) \leq \frac{1}{n_m} \sum_{k=1}^{n_m} U^k f(x_i) \leq \frac{C}{1-a}$$

for all $m \in \mathbb{N}$. Hence

$$\mu(f_R) \leq \frac{C}{1-a} \text{ for all } R > 0.$$

Therefore, by the Monotone Convergence Theorem,

$$\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) = \mu(f) \leq \frac{C}{1-a}.$$

□

Proposition 2 *Suppose $\mathcal{M}$ is a CMS such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Then $ES(u)$ is nonempty.*

Proof. By Proposition 1, the CMS has an invariant Borel probability measure μ such that

$$\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty \text{ for all } x_i \in K_i, i = 1, \dots, N.$$

Furthermore, by Lemma 6 (ii),

$$M(Y) = 1,$$

where M is the generalized Markov measure associated with $\mathcal{M}$ and μ . Hence, by Proposition 1 in [20], M is an equilibrium state for u . □

Lemma 7 *Let $\Lambda \in ES(u)$ and $\tilde{\Lambda} \in P_S(\Sigma)$ such that $\tilde{\Lambda}$ is absolutely continuous with respect to Λ . Then $\tilde{\Lambda} \in ES(u)$.*

Proof. Let ψ be the Radon-Nikodym density of $\tilde{\Lambda}$ with respect to Λ . By the shift invariance of $\tilde{\Lambda}$ and Λ , $\psi \circ S = \psi$. This implies that

$$E_\Lambda(\psi|\mathcal{F}) = E_\Lambda(\psi|\mathcal{F}) \circ S \text{ } \Lambda\text{-a.e..}$$

Let $A \in \mathcal{F}$ and $e \in E$. Then

$$\begin{aligned} \int_A 1_{1[e]} d\tilde{\Lambda} &= \int_A 1_{1[e]} \psi \circ S d\Lambda = \int_{S(A)} 1_{0[e]} E_\Lambda(\psi|\mathcal{F}) d\Lambda \\ &= \int_A 1_{1[e]} E_\Lambda(\psi|\mathcal{F}) d\Lambda = \int_A E_\Lambda(1_{1[e]}|\mathcal{F}) E_\Lambda(\psi|\mathcal{F}) d\Lambda \\ &= \int_A E_\Lambda(1_{1[e]}|\mathcal{F}) \psi d\Lambda = \int_A E_\Lambda(1_{1[e]}|\mathcal{F}) d\tilde{\Lambda}. \end{aligned}$$

Hence

$$E_{\tilde{\Lambda}}(1_{1[e]}|\mathcal{F}) = E_{\Lambda}(1_{1[e]}|\mathcal{F}) \quad \tilde{\Lambda}\text{-a.e..}$$

Since Λ is an equilibrium state for u ,

$$\sum_{e \in E} 1_{1[e]} \log E_{\Lambda}(1_{1[e]}|\mathcal{F}) = u \quad \Lambda\text{-a.e.,}$$

by Lemma 5 in [20]. Therefore, by the absolute continuity of $\tilde{\Lambda}$ with respect to Λ ,

$$\sum_{e \in E} 1_{1[e]} \log E_{\tilde{\Lambda}}(1_{1[e]}|\mathcal{F}) = u \quad \tilde{\Lambda}\text{-a.e..}$$

Therefore,

$$\begin{aligned} h_{\tilde{\Lambda}}(S) &= - \sum_{e \in E} \int E_{\tilde{\Lambda}}(1_{1[e]}|\mathcal{F}) \log E_{\tilde{\Lambda}}(1_{1[e]}|\mathcal{F}) \, d\tilde{\Lambda} \\ &= - \int \sum_{e \in E} 1_{1[e]} \log E_{\tilde{\Lambda}}(1_{1[e]}|\mathcal{F}) \, d\tilde{\Lambda} \\ &= - \int u \, d\tilde{\Lambda}. \end{aligned}$$

Since $P(u) = 0$ (see e.g. [20]), it follows that $\tilde{\Lambda} \in ES(u)$. $\square$

Proposition 3 *Suppose $\mathcal{M}$ is CMS such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Then there exists an ergodic equilibrium state Λ for the energy function u such that Λ is absolutely continuous with respect to M where M is a generalized Markov measure associated with the CMS and its invariant Borel probability measure μ with the property that $\sum_{i=1}^N \int_{K_i} d(x, x_i) \, d\mu(x) < \infty$ for all $x_i \in K_i$, $i = 1, \dots, N$.*

Proof. By Proposition 1, there exists an invariant Borel probability measure μ of the CMS such that $\sum_{i=1}^N \int_{K_i} d(x, x_i) \, d\mu(x) < \infty$ for all $x_i \in K_i$, $i = 1, \dots, N$. By Proposition 1 in [20], M is an equilibrium state for u . Since Σ is a compact metric space, it is well known that $P_S(\Sigma)$ is a compact (in the weak* topology) convex space. By the Krein-Milman Theorem, $P_S(\Sigma)$ is a closed convex hull of its extreme points, which are exactly the ergodic measures (see e.g. Theorem 6.10 in [15]). Furthermore, the ergodic measures can be characterized as the minimal elements of the set $P_S(\Sigma)$ with respect to the partial order given by the absolute continuity relation (see e.g. [9], Lemma 2.2.2) (to be precise, the absolute continuity relation is a partial order on the equivalence classes consisting of equivalent measures, but every equivalence class containing an ergodic measure consists of a single element). Therefore, there exists an ergodic $\tilde{\Lambda} \in P_S(\Sigma)$ such that $\tilde{\Lambda}$ is absolutely continuous with respect to M . Thus, by Lemma 7, $\tilde{\Lambda} \in ES(u)$. $\square$

Theorem 3 *Suppose CMS $\mathcal{M}$ has a unique invariant Borel probability measure μ and $M(Y) = 1$, where M is the associated generalized Markov measure. Then the following hold.*

- (i) M is a unique equilibrium state for the energy function u ,
- (ii) $P(u) = 0$,
- (iii) $F(M) = \mu$,
- (iv) $h_M(S) = -\sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e \, d\mu$.

For a proof see [20].

2.3 Uniqueness and empiricalness of the invariant probability measure

Now, we are going to show that an irreducible CMS with probabilities with square summable variation has a unique invariant probability measure and it can be obtained empirically.

Before we move to the main theorem, we need to clear up some technical details.

Let $\nu \in P(K)$. Since $x \mapsto P_x(Q)$ is Borel measurable for all $Q \in \mathcal{B}(\Sigma^+)$ (see e.g. Lemma 1 in [18]), we can define

$$\tilde{\phi}(\nu)(A \times Q) := \int_A P_x(Q) d\nu(x)$$

for all $A \in \mathcal{B}(K)$ and $Q \in \mathcal{B}(\Sigma^+)$. Then $\tilde{\phi}(\nu)$ extends uniquely to a Borel probability measure on $K \times \Sigma^+$ with

$$\tilde{\phi}(\nu)(\Omega) = \int P_x(\{\sigma \in \Sigma^+ : (x, \sigma) \in \Omega\}) d\nu(x)$$

for all $\Omega \in \mathcal{B}(K \times \Sigma^+)$. Note that the set of all $\Omega \subset K \times \Sigma^+$ for which the integrand in the above is measurable forms a Dynkin system which contains all rectangles. Therefore, it is measurable for all $\Omega \in \mathcal{B}(K \times \Sigma^+)$.

Now, consider the following map

$$\begin{aligned} \xi : \Sigma &\longrightarrow K \times \Sigma^+ \\ \sigma &\longmapsto (F(\sigma), (\sigma_1, \sigma_2, \dots)). \end{aligned}$$

Lemma 8 *Suppose $\mathcal{M}$ is a CMS with an invariant Borel probability measure μ such that $\sum_{i=1}^N \int_{K_i} d(x, x_i) \, d\mu(x) < \infty$ for all $x_i \in K_i$, $i = 1, \dots, N$, and $M(Y) = 1$, where M is the generalized Markov measure associated with $\mathcal{M}$ and μ . Let $\Lambda \in P_S(\Sigma)$ be absolutely continuous with respect to M . Then*

$$\xi(\Lambda) = \tilde{\phi}(F(\Lambda)).$$

Proof. We only need to check that

$$\xi(\Lambda) (A \times_1 [e_1, \dots, e_n]^+) = \tilde{\phi}(F(\Lambda)) (A \times_1 [e_1, \dots, e_n]^+)$$

for all cylinder sets ${}_1[e_1, \dots, e_n]^+ \subset \Sigma^+$ and $A \in \mathcal{B}(K)$. For such sets

$$\begin{aligned} \xi(\Lambda) (A \times_1 [e_1, \dots, e_n]^+) &= \Lambda (F^{-1}(A) \cap {}_1[e_1, \dots, e_n]) \\ &= \int_{F^{-1}(A)} 1_{{}_1[e_1, \dots, e_n]} d\Lambda \end{aligned}$$

where ${}_1[e_1, \dots, e_n] \subset \Sigma$ is the pre-image of ${}_1[e_1, \dots, e_n]^+$ under the natural projection. Recall that $F^{-1}(A) \in \mathcal{F}$. Therefore, by Lemma 6 in [20],

$$E_M(1_{{}_1[e]}|\mathcal{F}) = p_e \circ F \text{ } M\text{-a.e.}$$

Since Λ is absolutely continuous with respect to M ,

$$E_\Lambda(1_{{}_1[e]}|\mathcal{F}) = p_e \circ F \text{ } \Lambda\text{-a.e.}, \quad (5)$$

analogously as in the proof of Lemma 7. This implies, by the shift invariance of Λ and the pull-out property of the conditional expectation, that

$$E_\Lambda(1_{{}_1[e_1, \dots, e_n]}|\mathcal{F}) = P_{F(\sigma)} ({}_1[e_1, \dots, e_n]^+) \text{ } \Lambda\text{-a.e.}$$

Therefore,

$$\begin{aligned} \int_{F^{-1}(A)} 1_{{}_1[e_1, \dots, e_n]} d\Lambda &= \int_{F^{-1}(A)} P_{F(\sigma)} ({}_1[e_1, \dots, e_n]^+) d\Lambda(\sigma) \\ &= \int_A P_x ({}_1[e_1, \dots, e_n]^+) dF(\Lambda)(x), \end{aligned}$$

as desired. $\square$

Theorem 4 *Suppose $\mathcal{M}$ is an irreducible CMS such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Then the following hold.*

- (i) *The CMS has a unique invariant Borel probability measure μ .*
- (ii) *Let $f_e : K \rightarrow [-\infty, +\infty]$ be Borel measurable such that $f_e|_{K_{i(e)}}$ is bounded and uniformly continuous for all $e \in E$. Then, for every $x \in K$,*

$$\frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) \rightarrow \sum_{e \in E_{K_{i(e)}}} \int p_e f_e d\mu \text{ for } P_x\text{-a.e. } \sigma \in \Sigma^+.$$

Proof. By Proposition 3, u has an ergodic equilibrium state Λ which is absolutely continuous with respect to M where M is the generalized Markov measure

associated with the CMS and its invariant Borel probability measure μ with the property that $\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$ for all $x_i \in K_i$, $i = 1, \dots, N$. This implies that $\Lambda(Y) = 1$. Set

$$v(\sigma) = \begin{cases} \log f_{\sigma_1} \circ F(\sigma) & \text{if } \sigma \in Y \\ -\infty & \text{if } \sigma \in \Sigma \setminus Y. \end{cases} \quad (6)$$

Then, by Birkhoff's Ergodic Theorem,

$$\frac{1}{n} \sum_{k=0}^{n-1} v \circ S^k \rightarrow \int v d\Lambda \quad \Lambda\text{-a.e..}$$

Since $F \circ S^k(\sigma) = w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(F(\sigma))$ for all $\sigma \in Y$, we can assume, without loss of generality, that

$$\frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(F(\sigma)) \rightarrow \int v d\Lambda \quad \text{for all } \sigma \in Y.$$

By (5),

$$\begin{aligned} \int v d\Lambda &= \lim_{n \rightarrow \infty} \sum_{e \in E} \int 1_{[e]} n \wedge f_e \circ F d\Lambda = \lim_{n \rightarrow \infty} \sum_{e \in E} \int p_e \circ F n \wedge f_e \circ F d\Lambda \\ &= \lim_{n \rightarrow \infty} \sum_{e \in E_{K_{i(e)}}} \int p_e n \wedge f_e dF(\Lambda) = \sum_{e \in E_{K_{i(e)}}} \int p_e f_e dF(\Lambda). \end{aligned}$$

Now, applying the map ξ , we get

$$\frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) \rightarrow \sum_{e \in E_{K_{i(e)}}} \int p_e f_e dF(\Lambda) \quad \text{for all } (x, \sigma) \in \xi(Y).$$

By Lemma 8, $\tilde{\phi}(F(\Lambda))(\xi(Y)) = \xi(\Lambda)(\xi(Y)) = \Lambda(\xi^{-1}(\xi(Y))) \geq \Lambda(Y) = 1$. Since $\tilde{\phi}(F(\Lambda))$ is a probability measure,

$$\begin{aligned} 1 &= \tilde{\phi}(F(\Lambda))(\xi(Y)) = \int P_x(\{\sigma \in \Sigma^+ : (x, \sigma) \in \xi(Y)\}) dF(\Lambda)(x) \\ &= \sum_{i=1}^N \int_{K_i} P_x(\{\sigma \in \Sigma^+ : (x, \sigma) \in \xi(Y)\}) dF(\Lambda)(x). \end{aligned}$$

Furthermore, for each $i = 1, \dots, N$, there exists $x_i \in K_i$ such that

$$\begin{aligned} &\int_{K_i} P_x(\{\sigma \in \Sigma^+ : (x, \sigma) \in \xi(Y)\}) dF(\Lambda)(x) \\ &= P_{x_i}(\{\sigma \in \Sigma^+ : (x_i, \sigma) \in \xi(Y)\}) F(\Lambda)(K_i). \end{aligned}$$

Set

$$Q_i := \{\sigma \in \Sigma^+ : (x_i, \sigma) \in \xi(Y)\}.$$

Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x_i) = \sum_{e \in E_{K_i(e)}} \int p_e f_e dF(\Lambda) \text{ for all } \sigma \in Q_i,$$

and

$$\sum_{i=1}^N P_{x_i}(Q_i) F(\Lambda)(K_i) = 1.$$

Note that, since Λ is an equilibrium state for u , $U^*F(\Lambda) = F(\Lambda)$ (e.g. Proposition 1 in [20]). Thus, by the irreducibility of the CMS $F(\Lambda)(K_i) > 0$ for all $i = 1, \dots, N$ (see Lemma 1 in [16]). This implies that $P_{x_i}(Q_i) = 1$ for all $i = 1, \dots, N$, as $F(\Lambda)$ and P_{x_i} are probability measures. Now, fix $x \in K_i$ for some $i \in \{1, \dots, N\}$. Since P_x is absolutely continuous with respect to P_{x_i} , $P_x(Q_i) = 1$. Furthermore, the contractiveness on average condition (1) implies that

$$\int d(w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x), w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x_i)) dP_x \leq a^k d(x, x_i).$$

Therefore, it follows, by the Borel-Cantelli argument, that

$$d(w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x), w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x_i)) \rightarrow 0 \text{ } P_x\text{-a.e..}$$

As each $f_e|_{K_i(e)}$ is uniformly continuous, we conclude that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) = \sum_{e \in E_{K_i(e)}} \int p_e f_e dF(\Lambda) \text{ } P_x\text{-a.e..}$$

Thus, the claim in (ii) will follow if we show that $F(\Lambda)$ is the unique invariant Borel probability measure of the CMS.

Let $g \in C_B(K)$ and set $f_e = g$ for all $e \in E$. Then the integration with respect to P_x and Lebesgue's dominated convergence theorem imply that

$$\frac{1}{n} \sum_{k=0}^{n-1} U^k g(x) \rightarrow \int g dF(\Lambda) \text{ for all } g \in C_B(K).$$

Since x was arbitrary, again an integration and Lebesgue's dominated convergence theorem imply that

$$\frac{1}{n} \sum_{k=0}^{n-1} U^{*k} \nu \xrightarrow{w^*} F(\Lambda) \text{ for all } \nu \in P(K). \quad (7)$$

Suppose $U^*\lambda = \lambda$ for some $\lambda \in P(K)$. Then

$$\frac{1}{n} \sum_{k=0}^{n-1} U^{*k} \lambda = \lambda \text{ for all } n \in \mathbb{N}.$$

Therefore, $F(\Lambda) = \lambda$. Hence, $F(\Lambda)$ is a unique invariant probability measure of the CMS, i.e. $F(\Lambda) = \mu$. $\square$

The following corollary generalizes corresponding results from [1], [6], [16] and [19].

Corollary 2 *Suppose $\mathcal{M}$ is an irreducible CMS such that each $p_e|_{K_{i(e)}}$ has a square summable variation. Then the following hold.*

- (i) *The CMS has a unique invariant Borel probability measure μ .*
- (ii) *Let $f_e : K \rightarrow [-\infty, +\infty]$ be Borel measurable such that $f_e|_{K_{i(e)}}$ is bounded and uniformly continuous for all $e \in E$. Then, for every $x \in K$,*

$$\frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) \rightarrow \sum_{e \in E_{K_{i(e)}}} \int p_e f_e d\mu \text{ for } P_x\text{-a.e. } \sigma \in \Sigma^+.$$

Proof. By Lemma 2, the measures P_x and P_y are absolutely continuous with respect to each other for all $x, y \in K_i$, $i = 1, \dots, N$. Therefore, the claim follows by Theorem 4. $\square$

Corollary 3 *Suppose $\mathcal{M}$ is an irreducible CMS such that each $p_e|_{K_{i(e)}}$ has a square summable variation. Then the following hold.*

- (i) *The generalized Markov measure M is a unique equilibrium state for the energy function u ,*
- (ii) *$P(u) = 0$,*
- (iii) *$F(M) = \mu$,*
- (iv) *$h_M(S) = -\sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e d\mu$.*

Proof. Since the CMS has a unique invariant Borel probability measure μ (Corollary 2) and $M(Y) = 1$ (Lemma 6 (ii)), the claim follows by Theorem 3 and Theorem 4. $\square$

Finally, we would like to give an application of Theorem 4 (iv) which allows an empirical calculation of Kolmogorov-Sinai entropy $h_M(S)$ of the generalized Markov shift associated with a given CMS without knowing anything about its invariant measure.

Corollary 4 *Suppose $\mathcal{M}$ is an irreducible CMS such that P_x is absolutely continuous with respect to P_y for all $x, y \in K_i$, $i = 1, \dots, N$. Then for every $x \in K$*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log P_x([\sigma_1, \dots, \sigma_n]) = -h_M(S) \text{ for } P_x\text{-a.e. } \sigma \in \Sigma^+.$$

Proof. By Theorem 4 (i) and Theorem 3 (iv),

$$h_M(S) = - \int \sum_{e \in E} p_e \log p_e d\mu.$$

Set $f_e := \log p_e$ for all $e \in E$. Then, by Theorem 4 (ii), for every $x \in K$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log[p_{\sigma_1}(x)p_{\sigma_2} \circ w_{\sigma_1}(x) \dots p_{\sigma_n} \circ w_{\sigma_{n-1}} \circ \dots \circ w_{\sigma_1}(x)] = -h_M(S)$$

for P_x -a.e. $\sigma \in \Sigma^+$, as desired. $\square$

Remark 4 The author would like to point out that a similar entropy formula as that proved in Theorem 3 (iv) plays a central role in the recent book of Wojciech Ślomeczynski [13].

Remark 5 This paper contributes more to the mystery of an ergodic classification of the generalized Markov shifts associated with aperiodic CMSs. So far, we know only that they are strongly mixing if the restrictions of the probability functions on their vertex sets are Dini-continuous [19] and they are weak Bernoulli if in addition all maps are contractive (the latter is not difficult to see by reducing it to the corresponding result of P. Walters for the natural extension of g -measures [14] since $g := \exp u|_{\Sigma_G}$ is Dini-continuous in this case). H. Brebee [2] has shown the very weak Bernoulli property of the natural extension of g -measures on a full shift under a continuity condition on g which is weaker than the Dini-continuity and not comparable with the square summability of variation (see [10]). From this paper, we only can see that an aperiodic generalized Markov shift is ergodic if the restrictions of the probability functions on their vertex sets have a square summable variation (Corollary 3 and Lemma 3).

Acknowledgements

I would like to thank Barry Ridge, Wang Yang and Zhenkun Wang for their help in the production of this paper.

References

- [1] M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo, Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **24** (1988) 367-394.
- [2] H. Brebee, Chains with Infinite Connections: Uniqueness and Markov Representation, *Probab. Th. Rel. Fields* **76** (1987), 243-253.
- [3] N. Berger, Ch. Hoffman, V. Sidoravicius, Nonuniqueness for specifications in $\ell^{2+\epsilon}$, preprint *math.PR/0312344*.

- [4] L. Breiman, *Probability*, Addison-Wesley Publishing Company, Reading, Mass.-London-Don Mills, Ont. (1968).
- [5] J. L. Doob, *Measure Theory*, Spriger-Verlag (1993).
- [6] J. H. Elton, An ergodic theorem for iterated maps, *Ergod. Th. & Dynam. Sys.* **7** (1987) 481-488.
- [7] R. Isaac, Markov processes and unique stationary probability measures, *Pacific J. Math.* **12** (1962) 273-286.
- [8] M. Keane, Strongly Mixing g -Measures, *Inventiones math.* **16** (1972) 309-324.
- [9] G. Keller, *Equilibrium States in Ergodic Theory*, Student Texts **42** LMS, Cambridge University Press (1998).
- [10] A. Johansson and A. Öberg, Square summability of variations of g -functions and uniqueness of g -measures, *Math. Res. Lett.* **10** (2003), 587-601.
- [11] F. Ledrappier, Principe variationnel et systèmes dynamiques symboliques, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **30** (1974), 185-202.
- [12] R. D. Mauldin and S. C. Williams, Hausdorff dimension in graph directed constructions, *Tran. AMS Vol 309 (1988)* 811-829.
- [13] W. Ślomczyński, *Dynamical entropy, Markov operators, and iterated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).
- [14] P. Walters, Ruelle's Operator Theorem and g -measures, *Tran. AMS* **214** (1975) 375-387.
- [15] P. Walters, *An introduction to Ergodic Theory*, Springer-Verlag, (1982).
- [16] I. Werner, Contractive Markov systems, *J. London Math. Soc.* **71** (2005) 236-258.
- [17] I. Werner, Contractive Markov System with Constant Probabilities, *J. Theoret. Prob.* **18** (2005), no. 2, 469-479.
- [18] I. Werner, Coding map for a contractive Markov system, *Math. Proc. Camb. Phil. Soc.* to appear **140** (2006), *preprint math.PR/0504247*.
- [19] I. Werner, Ergodic theorem for a contractive Markov system, *Nonlinearity* **17** (2004) 2303-2313.
- [20] I. Werner, The generalized Markov measure as an equilibrium state, *Nonlinearity* **18** (2005) 2261-2274.